

Tooth whitening: the last 25 years

Linda Greenwall investigates tooth whitening and how it has become a valued treatment service for patients with a vast body of scientific literature behind it

The introduction of contemporary tooth whitening techniques was launched in 1989 with a research paper called 'Nightguard vital bleaching'. The authors, Van Haywood and Harald Heymann, described the use of a bleaching tray as a vehicle to place the whitening gel into the mouth for better retention and to give better long lasting and predictable results.

Further research by the authors showed the effectiveness, efficacy, predictability, longevity and reduction in side effects.

It has now been 27 years since that paper was published and this article will review how, in just a quarter of a century, tooth whitening has become a valued treatment service for patients with a vast body of scientific literature behind it.

Millions of people around the world have benefitted from the use of the tooth whitening materials and techniques over the last 25 years and its popularity continues to grow (Greenwall, 2001).

Although tooth whitening techniques were popular in the 1880s – using strong concentrations of hydrogen peroxide and a bleaching lamp – modern techniques focus on the use of a bleaching tray and applying products in the tray at home.

It was Bill Klausmier, an American orthodontist, who started using the technique in 1968 to help reduce the swelling on the gingivae post-orthodontic treatment. He



Education aims and objectives

This article aims to explain how tooth whitening has developed over the last 25 years.

Expected outcomes

Correctly answering the questions on page 66, worth one hour of verifiable CPD, will demonstrate that the reader understands how tooth whitening has become a valued treatment service for patients with a vast body of scientific literature behind it.



Figure 1: An advanced whitening case with a diagnosis of fluorosis. The patient had direct composite bondings placed over the upper central incisor teeth to mask the discolouration, which were removed prior to whitening. Home bleaching was undertaken using 10% carbamide peroxide in a bleaching tray



Figure 2: Result after whitening with carbamide peroxide for eight to 10 weeks

advised his patients to use peroxy mouthwash into the retainer to reduce the gingival hyperplasia (Haywood, 1991a). At the six-month recalls he noted that not only were the gingivae of the patient significantly better, but the teeth were also whiter.

This chance finding, similar to the invention of penicillin, has allowed many patients to benefit from these techniques. After using this technique for 40 years, he reported that nobody needed a root canal treatment, nor broke nor damaged a tooth following the use of peroxide in the tray. He advised his colleagues in a local study group to use this technique and subsequently passed it onto Dr Van Haywood who began research on the technique.

Early research

Early research focused on whether the whitening products were safe and effective. Professor Yiming Li from Loma Linda University has devoted the last 20 years of his

research life investigating the safety of hydrogen peroxide. He has concluded that it is safe to use as a whitening agent in the oral cavity, as long as the products used are supervised and monitored by the dentist and the dental team (Li, Greenwall, 2013).

There was an explosion of research conducted on whitening in the early 1990s, and there are now thousands of articles published on all aspects of whitening.

There was extensive research conducted on the side effect of sensitivity, which occurs in up to 85% of patients. Research was conducted as to how and why the sensitivity occurred and how best to treat it.

It was discovered that whitening gel penetrates the tooth within five to 15 minutes of application. It is therefore essential to assess the pulps of all the teeth to make sure they are healthy prior to undertaking any whitening procedure. If whitening was to be undertaken in the presence of a non-vital area, the area

Dr Linda Greenwall is a specialist in prosthodontics and restorative dentistry, and principal of the award-winning Heath Dental Care in the heart of London's Hampstead. An international authority on tooth whitening, she is the founder and current chair of the British Dental Bleaching Society, and the driving force behind the international charity, the Dental Wellness Trust. Dr Greenwall is editor-in-chief of the journal *Aesthetic Dentistry Today*, and has written several textbooks, including the recently updated second edition of *Bleaching Techniques in Restorative Dentistry*.

How has tooth whitening changed over time?		
Patient factors	Material factors	Technique factors
Expectations increased	Two-week tray use now	Take home gels now fifth generation formulation
Seeking whiter teeth	Extended tray use	Introduction of soothers, potassium nitrate and fluoride and acp
Philosophy of perfection	Changes in tray designs	Concentrations of materials have changed
More difficult discolourations	Used with aligners	Power gels have changed
No age restrictions for older age patients	Therapeutic uses now introduced	Is light essential?
No age limit for under 18	Whitening strips introduced	Lights/no light?
Whitening maintenance		Heat/no heat?
Whitening for life		Lasers
Bleachorexia/bleachoholic		Ozone

Table 1: Some of the changes that have occurred in tooth whitening over the last 25 years

will flare up and need a root canal treatment. It is best to plan a root canal treatment to be undertaken prior to starting any whitening treatment.

Methods of whitening

There are two basic techniques for home bleaching, depending on the products:

- Night-time use, using a carbamide peroxide material
- Day-time use, using a hydrogen peroxide only material.

How have things changed?

Changes in the whitening legislation were introduced in 2011. The Cosmetic Products (Safety) (Amendment) Regulations 2012 (implementing Directive 2011/84/EU, which amends Directive 76/768/EEC) came into force on 31 October 2012.

The legislation specifies that products containing or releasing between 0.1% and six per cent hydrogen peroxide cannot be used on any person under the age of 18, except where such use is intended wholly for the purpose of treating or preventing disease.

Products containing or releasing less than 0.1% of hydrogen peroxide, including mouth rinse, toothpaste and tooth whitening or bleaching products, are safe and will continue to be freely available on the market.

Tooth whitening or bleaching products containing or releasing between 0.1% and six per cent of hydrogen peroxide may be used provided an appropriate clinical examination is carried out in order to ensure that there are no risk factors and any other oral pathology is ruled out.

The exposure to the whitening products should be limited to ensure that they are only

used as intended in terms of frequency and duration of application.

Whitening products should be clearly labelled for their intended purpose of tooth whitening. The products should not be directly freely available to the consumer. Products may only be purchased through a dental practice and tooth whitening products containing or releasing between 0.1% and six per cent hydrogen peroxide may only be sold to dental practitioners.

For each cycle of use, the first use must be carried out by a dental practitioner or under their direct supervision if an equivalent level of safety is ensured. After the first cycle of use, the dental practitioner may give the product to the consumer to complete the cycle of use.

Concentrations exceeding six per cent of hydrogen peroxide remain prohibited unless wholly for the purpose of the treatment or prevention of disease. It is essential that dentists abide by the legislation, as Trading Standards can prosecute them if they use higher than six per cent hydrogen peroxide on a patient or if they intend to supply a patient with whitening gel stronger than six per cent hydrogen peroxide.

Bleaching single dark teeth

The treatment options for both vital and non-vital teeth have changed over the last 25 years with the introduction of whitening techniques.

These days a sectional whitening tray can be used to whiten a single dark tooth, whether vital or non-vital. It is essential to whiten the dark tooth first in order to establish the whitening potential and possibility. If the whitening of the dark tooth doesn't start first, the rest of the teeth will whiten quicker and the contrast of the dark tooth will look worse.

Vital single dark teeth

When the patient receives mild trauma to an anterior tooth, the tooth tries to repair and heal itself by laying down extra secondary and tertiary dentine in the pulp chamber. The trauma results in bleeding into the pulp chamber and pulp canal. The blood products reorganise and break down into iron. The discolouration, which is a result of the bleeding and the formation of secondary and tertiary dentine, can be detected when the tooth is a different colour to its neighbour by about one or two shades only.

In the past it was thought that these teeth, which are diagnosed as having calcific metamorphosis (Haywood, 2010), need to have root canal treatments. However, this is not true – they need only be whitened.

Non-vital teeth

The techniques for non-vital bleaching have also evolved with the banning of sodium perborate by the scientific committee in Europe, which was concerned about the fetotoxic and cytotoxic effects.

The standard technique, which was called the Walking Bleach technique and first described by Nutting and Poe in 1965, was advocated by using sodium perborate mixed with 35% hydrogen peroxide.

The two products together act synergistically and create the equivalent of 50% hydrogen peroxide, which is too caustic for a root canal considering that it may be affected by trauma previously. High concentrations of hydrogen peroxide have been banned in Europe and it may only be possible to use six per cent hydrogen peroxide sealed into a root canal.

In addition, the strong concentrations of hydrogen peroxide in combination with previous trauma to the tooth may result in cervical resorption (Cvek, Lindwall, 1985; Hierthersay 1999), which has been extensively described in the literature.

There are new modifications for this treatment, which involve the use of a bleaching tray and a segmental bleaching tray. These days, 16% carbamide peroxide is sealed into the root canal and the patient uses the bleaching tray to whiten the external surface of the tooth, meaning that the tooth is effectively whitened from the inside and the outside with the same technique.

Dr Willie Liebenberg described a modification in 1997, in which he advocated leaving the access cavity of the non-vital tooth open so that the patient could apply the whitening syringe into the access cavity every two hours thereby whitening the tooth over

involves regular professional oral prophylaxis treatment, the use of whitening toothpaste and sometimes a reduction in food and drink that cause staining.

The rise of bleachorexia

Over the last 25 years it has been possible to whiten the teeth beyond the original Vita classic shade guides.

New shade guides have been developed to match the new shades of white, and porcelain and composite shades have been introduced onto the market to be able to restore these teeth to the new whiter shade.

Some patients have developed a syndrome where they continually seek whiter and whiter teeth. This is also associated with body dysmorphic disorder and low self-esteem. The term has been described as bleachorexia (Kelleher, 2014), with patients referred to as being a bleachoholic.

It is essential that dentists know how to recognise the syndromes and not be seduced into undertaking unnecessary whitening treatment. These patients can usually be detected early, as their teeth are whiter than their sclera of their eyes, which is often used as a measurement of the whiteness that can be achieved.

It is essential that when whitening treatment is completed the patient is told that no further whitening is necessary at that point. Each whitening cycle will require the dentist to reassess the patient to assess whether further whitening is necessary.

Conclusion

The explosion in tooth whitening techniques research over the last 25 years has demonstrated

that the whitening products are safe, effective, predictable to use and have many benefits for patients. Side effects such as sensitivity can be managed well as the newer generations of whitening products now contain extra soothers such as potassium nitrate, fluoride and amorphous calcium phosphate. Patients have benefitted from their new shade of white teeth and these treatments have improved smiles in a natural and non-invasive way.

Further clarification in the whitening legislation for under 18s (Kelleher, 2014) is


being investigated by the CED in Europe, and these treatments may be able to be used for under 18s, provided they are being used for the treatment of disease, which they are.

It is expected that the amount of professional tooth whitening materials available will increase for patients to use under the direction of dentists. What's more, there are many new innovative whitening products that are being brought onto the market for patients to continue to enjoy a whiter, brighter smile.

Care to comment? @AesDenToday

References

- Cvek M, Lindwall A M (1985) External root resorption following bleaching of pulpless teeth with oxygen peroxide. *Endod Dent Traumatol* 1: 56-60
- Greenwall LH (2001) *Bleaching techniques in Restorative Dentistry*. First edition. Martin Dunitz, aka Taylor and Francis, London
- Greenwall LH (2009) Treatment considerations for bleaching and bonding white lesions in the anterior dentition. *Alpha Omegan* 102(4): 121-7
- Haywood VB (1991a) Overview and status of mouthguard bleaching. *J Esthet Dent* 3(5): 157-61
- Haywood VB (1991b) Nightguard Vital Bleaching, a history and products update: Part 1. *Esthet Dent Update* 2(4): 63-6
- Haywood VB (1991c) Nightguard Vital Bleaching, a history and products update: Part 2. *Esthet Dent Update* 2(5): 82-5
- Haywood VB, DiAngelis AJ (2010) Bleaching the Single Dark Tooth. Changing the color of just one anterior tooth presents unique challenges. *Inside Dentistry* 42-52
- Haywood VB (2010) Orthodontic caries control and bleaching. *Inside Dentistry* 6(4): 36-50
- Heithersay GS (1999) Invasive cervical resorption: an analysis of potential predisposing factors. *Quintessence Int* 30: 83-95
- Kelleher M (2014) The law is an ass: ethical and legal issues surrounding the bleaching of young patients' discoloured teeth. *Faculty Dental Journal* 5(2): 56-67
- Kelleher M, Djemal S, Ahmed S, Al-Khayatt, Ray-Chaudhuri AJ, Briggs P, Porter RW (2011) Bleaching and Bonding for the older patient. *Dental Update* 38: 294-303
- Kelleher MGD, Roe F (1999) The safety-in- use of 10% carbamide peroxide (Opalescence) for bleaching teeth under the supervision of a dentist. *Br Dent J* 187: 190-194
- Li Y, Greenwall L (2013) Safety issues of tooth whitening using peroxide-based materials. *Br Dent J* 215(1): 29-34
- Liebenberg WH (1997) Intracoronal lightening of discoloured pulpless teeth: a modified walking bleach technique. *Quintessence Int* 28: 771-777
- Nutting E B, Poe G S (1967) Chemical bleaching of discoloured endodontically treated teeth. *Dent Clin North Am* 11: 655-622
- Zimmerli B, Jeger F, Lussi A (2010) Bleaching of nonvital teeth. A clinically relevant literature review. *Schweiz Monatsschr Zahnmed* 120(4): 306-20



Private Dentistry is an essential reference journal, promoting excellence to private dental practitioners with editorially independent articles by opinion leaders, covering the full range of clinical and management topics.

~ 12 issues per year, 24 verifiable CPD hours

~ Clinical excellence step-by-step guides

~ Private life articles written by those succeeding in private dentistry

~ Expert advice on how to market your practice, keep your team motivated and increase revenue

~ News, updates and advice on the key issues affecting private dentists

